

5171 Wilfong Road Memphis, TN 38134 USA

Telephone: 901-382-8716 Fax: 901-333-8223

Email: info@meridianlifescience.com www.MeridianLifeScience.com

CERTIFICATE OF ANALYSIS

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog #: C01795M **Lot #:** 57L33615

Description: MAb to SED

Monoclonal Antibody to Staphylococcus aureus Enterotoxin D (SED)

Specificity: Specific to *Staphylococcus aureus* enterotoxin D. No cross reaction to other types of *Staphylococcus aureus*

enterotoxin A, B, C, and E.

Clone: B1915M

Host Animal: Mouse. Hybridization of P3X63.Ag8.653 myeloma cells **Isotype:** IgG₁

with spleen cells from BALB/c mice.

Immunogen: Staphylococcus aureus Enterotoxin D

Source: Ascites

Format: Purified, Liquid

Purification: $\geq 95\%$ pure (SDS-PAGE). Protein A Chromatography

Concentration: 3.6 mg/mL (OD280nm, $E^{0.1\%} = 1.38$)

Buffer: PBS: 1.5 mM Monopotassium Phosphate (KH₂PO₄), 8.0 mM Disodium Phosphate (Na₂HPO₄), 3 mM

Potassium Chloride (KCl), 140 mM Sodium Chloride (NaCl), pH 7.4.

Preservative: 0.05% Sodium Azide

Applications: Suitable for use in sandwich ELISA. Each laboratory should determine an optimum working titer for use in

its particular application. Other applications have not been tested but use in such assays should not

necessarily be excluded.

Recommended antibody pair for sandwich immunoassay:

<u>Capture</u> <u>Detection</u> C01795M C01793M

Storage: Short term (up to 7 days) store at 2-8°C. Long term, aliquot and store at -20°C. Avoid multiple freeze/thaw

cycles.

Warning: This product contains sodium azide, which has been classified as Xn (Harmful), in European Directive

67/548/EEC in the concentration range of 0.1 - 1.0 %. When disposing of this reagent through lead or

copper plumbing, flush with copious volumes of water to prevent azide build-up in drains.

03 Dec 2015

Signature