

## **CERTIFICATE OF ANALYSIS**

Centrifuge before opening to ensure complete recovery of vial contents.		
C88205M	Lot #:	1L34407
MAb to Influenza A H5N1 Monoclonal Antibody to Influenza A H5N1 (Avian) Hemaggl	utinin	
Hemagglutinin protein from the H5N1 strain of avian influenza (Middle Region).		
Mouse	Isotype:	IgG
Ascites		
Synthetic peptide corresponding to 14 amino acids near the middle of the Hemagglutinin protein (Genbank No. AAT76166)		
Affinity Purified, Liquid		
Immunoaffinity chromatography		
1 mg/ml		
Not determined		
Phosphate Buffered Saline		
0.02% Sodium azide		
Suitable for use in ELISA. It will detect 10ng of free peptide at 1ug/ml. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.		
Store (up to 1 year) at 2–8°C.		
Refer to the appropriate Safety Data Sheet (SDS) for additional information.		
	Centrifuge before opening to ensure complete real C88205M MAb to Influenza A H5N1 Monoclonal Antibody to Influenza A H5N1 (Avian) Hemaggl Hemagglutinin protein from the H5N1 strain of avian influenza Mouse Ascites Synthetic peptide corresponding to 14 amino acids near the m (Genbank No. AAT76166) Affinity Purified, Liquid Immunoaffinity chromatography 1 mg/ml Not determined Phosphate Buffered Saline 0.02% Sodium azide Suitable for use in ELISA. It will detect 10ng of free peptide a optimum working titer for use in its particular application. Oth such assays should not necessarily be excluded. Store (up to 1 year) at 2–8°C.	Centrifuge before opening to ensure complete recovery of vial content   C88205M Lot #:   MAb to Influenza A H5N1 Monoclonal Antibody to Influenza A H5N1 (Avian) Hemagglutinin   Hemagglutinin protein from the H5N1 strain of avian influenza (Middle Region). Mouse   Mouse Isotype:   Ascites Synthetic peptide corresponding to 14 amino acids near the middle of the Hemagglutinin (Genbank No. AAT76166)   Affinity Purified, Liquid Immunoaffinity chromatography   1 mg/ml Not determined   Phosphate Buffered Saline 0.02% Sodium azide   Suitable for use in ELISA. It will detect 10ng of free peptide at 1ug/ml. Each laboratory soptimum working titer for use in its particular application. Other applications have not be such assays should not necessarily be excluded.   Store (up to 1 year) at 2–8°C. Refer to the appropriate Safety Data Sheet (SDS) for additional information.



5171 Wilfong Road Memphis, TN 38134 USA Telephone: 901-382-8716 Fax: 901-333-8223 Email: info@meridianlifescience.com www.MeridianLifeScience.com

Catalog #C88205M Page 2 of 2

## **References**:

The references listed below are for research purposes only.

- 1. Thompson, W.W., et al., (2003), "Mortality associated with influenza and respiratory syncytial virus in the United States", JAMA, **289**: 179–186.
- Alexander, D.J., (2000), "A review of avian influenza. Proceedings of the European Society for Veterinary Virology (ESVV) Symposium on Influenza Viruses of Wild and Domestic Animals", <u>Vet.</u> <u>Microbiol.</u>, 74: 3–13.
- 3. Shortridge, K.F., et al., (1998), "Characterization of avian H5N1 influenza viruses from poultry in Hong Kong", <u>Virol</u>. **252**: 331–342.
- 4. Buxton Bridges, C., et al., (2000), "Risk of influenza A (H5N1) infection among health care workers exposed to patients with influenza A (H5N1), Hong Kong", J. Inf. Dis., **181**: 344–348.
- Iwatsuki-Horimoto, K., et al., (2004), "The index influenza A virus subtype H5N1 isolated from a human in 1997 differs in its receptor-binding properties from a virulent avian influenza virus", <u>J. Gen. Virol.</u>, 85: 1001–1005.

09 April 2017 Date

Signature

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY