

CERTIFICATE OF ANALYSIS

Important Note:	Centrifuge before opening to ensure complete recovery of vial contents.		
Catalog #: Page 1 of 2	K74190G	Lot #:	11126322
Description:	Goat anti Apo E Goat Antibody to Human Apolipoprotein E (Apo E)	
Specificity:	Binds to Apo E. Recognizes isoforms E2, E3 and E	E4.	
Host Animal:	Goat		
Immunogen:	Purified human Apo E from human plasma.		
Format:	Affinity Purified, Liquid		
Purification:	Human Apo E-Sepharose Affinity Column		
Concentration:	1.1 mg/mL (OD280nm, $E^{0.1\%} = 1.35$)		
Buffer:	75 mM Sodium Phosphate, 75 mM Sodium Chloride, 0.5 mM EDTA, pH 7.2.		
Preservative:	0.02% Sodium Azide		
Applications:	Can be used for detection of Apo E in plasma and lipoproteins; for ELISA and Western Blot (1:200 – 1:80,000), enzyme conjugation and biotinylation. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.		
Storage:	Short-term store at 2-8° C. Long term store at -20°C. Avoid multiple freeze/thaw cycles.		
Safety Notes (s):	Refer to the appropriate Safety Data Sheet (SDS) for additional information.		



5171 Wilfong Road Memphis, TN 38134 USA Telephone: 901-382-8716 Fax: 901-333-8223 Email: info@meridianlifescience.com www.MeridianLifeScience.com

Catalog #K74190G Page 2 of 2

References:

The references listed below are for research purposes only:

- 1. Wassef, H., et al., (2004), "Synthesis and Secretion of ApoC-I and ApoE during Maturation of Human SW872 Liposarcoma Cells", J. Nutr., 134: 2935-2941.
- DeMattos, R.B., et al., (1999), "A Test of the Cytosolic Apolipoprotein E Hypothesis Fails to Detect the Escape of Apolipoprotein E from the Endocytic Pathway into the Cytosol and Shows that Direct Expression of Apolipoprotein E in the Cytosol is Cytotoxic", <u>The Journal of Neuroscience</u>, **19**(7): 2464-2473.
- 3. Hasty, A.H., et al., (1999), "Retroviral Gene Therapy in ApoE-Deficient Mice ApoE Expression in the Artery Wall Reduces Early Foam Cell Lesion Formation", <u>Circulation</u>, **99**: 2571-2576.
- 4. Kitchens, R.L., et al., (2003), "Acute inflammation and infection maintain circulating phospholipid levels and enhance lipopolysaccharide binding to plasma lipoproteins", <u>J. Lipid Res.</u>, **44**: 2339-2348.
- 5. Swift, L.L., et al., (2001), "A Recycling Pathway for Resecretion of Internalized Apolipoprotein E in Liver Cells", <u>The Journal of Biological Chemistry</u>, **276**(25): 22965-22970.
- 6. Wilhelmus, M.M.M., (2005), "Apolipoprotein E Genotype Regulates Amyloid-beta Cytotoxicity", <u>The</u> <u>Journal of Neuroscience</u>, **25**(14): 3621-3627.
- 7. Böttcher, A., et al., (2000), "Preparative free-solution isotachophoresis for separation of human plasma lipoproteins: apolipoprotein and lipid composition of HDL subfractions", J. Lipid Res., **41**: 905-915.

Quality Signature:

20 Sep 2022

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY

SeanKlin